East African Scholars Journal of Psychology and Behavioural Sciences
Abbreviated Key Title: EAS J Psychol Behav Sci
ISSN 2663-1865 (Print) | ISSN 2663-6751 (Online)
Published By East African Scholars Publisher, Kenya

Volume-8 | Issue-4 | Oct-Dec-2026 | DOI: https://doi.org/10.36349/easjpbs.2026.v08i04.002

OPEN ACCESS

Original Research Article

Assessement of Spatial and Temporal Distribution of Measles and Yellow Fever and Vaccine Coverage Gaps in North-Central Nigeria, 2020–2024: a Mixed-Methods Study

Vincent-Vera Okwuosa[1*], Abah Michael Idoko[1,3], Adamu Ishaku Akyala[1], Olasunkanmi Samson Coker[1,2], Ikenna Osemeka Okeke[1], Onojah Matthew[1]

[1]Department of Global Health and Infectious Disease Control Institute, Nasarawa State University, Keffi, Nasarawa State, Nigeria
[2]Department of Epidemiology and Disease Control Technology, Federal College of Veterinary and Medical Laboratory Technology (FCVMLT), Vom, Nigeria
[3]Department of Haematology, Federal College of Veterinary and Medical Laboratory Technology (FCVMLT), Vom, Nigeria

Article History
Received: 06.08.2026
Accepted: 29.09.2026
Published: 03.10.2026

Journal homepage:
https://www.easpublisher.com

Quick Response Code

Abstract: ***Background***: Measles and Yellow Fever are vaccine-preventable diseases with recurrent outbreaks in Nigeria. North-Central Nigeria experiences seasonal peaks but lacks granular LGA-level spatial-temporal analysis. ***Methods:*** Concurrent mixed-methods design. Quantitative: NCDC surveillance sitreps 2020-2024, DHS 2021, N=1,842 household questionnaires. Qualitative: 24 KIIs and 8 FGDs. Spatial analysis used ArcGIS 10.8. Temporal analysis used epi-curves and seasonal decomposition. Vaccine coverage compared to WHO 80% threshold. STATA 17 for statistics. ***Results:*** 2,620 confirmed measles cases, 0 confirmed Yellow Fever cases 2020-2024. Nasarawa, Niger, and Kogi had highest burden: 210, 186, and 175 cases. Strong seasonality: Jan-April peaks, March 2.7x baseline. No state met WHO 80% threshold: Plateau 52%, Nasarawa 61%, Benue 55%, Kwara 46%, Kogi 44%. 70.4% of confirmed measles cases were zero-dose children. 2022 and 2024 showed national surges. ***Conclusions***: Persistent clustering, dry-season seasonality, and sub-threshold coverage drive outbreaks. Pre-season LGA-targeted campaigns before January and routine immunization strengthening are critical for WHO 2030 elimination goals.

Keywords: Measles, Yellow Fever, Vaccine Coverage, Spatial Epidemiology, Nigeria, Outbreak, North-Central.

Copyright © 2026 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution **4.0 International License (CC BY-NC 4.0)** which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

1. BACKGROUND

Emerging infectious diseases continue to threaten global health security, particularly in low- and middle-income countries with fragile health systems. Among vaccine-preventable diseases, measles and Yellow Fever remain public health priorities due to high transmissibility and potential for explosive outbreaks [1, 2].

Globally, measles caused an estimated 136,200 deaths in 2022, a 43% increase from 2021, driven by COVID-19-related immunization disruptions. The WHO Measles and Rubella Strategic Framework 2021-2030 targets elimination in 5 WHO regions by 2030. For Yellow Fever, the Eliminate Yellow Fever Epidemics (EYE) Strategy 2017-2026 aims to protect 1 billion people and eliminate outbreaks by 2026, with an 80% coverage threshold in at-risk areas [3-5].

In Nigeria, both diseases co-circulate in the North-Central zone. The region includes Plateau, Nasarawa, Benue, Niger, Kogi, Kwara, and the FCT, with a combined population of 24.8 million in 2024. The area is characterized by: 1) High population mobility along Abuja-Keffi and Minna-Suleja corridors with 122,000 and 95,500 monthly travelers; 2) Ecological suitability for _Aedes aegypti_ with forest cover 22-47% and Breteau Index >30 in sentinel sites; 3) Distinct dry season Jan-April that promotes measles transmission via low humidity and increased social gatherings [6-9].

***Corresponding Author:** Vincent-Vera Okwuosa
Department of Global Health and Infectious Disease Control Institute, Nasarawa State University, Keffi, Nasarawa State, Nigeria

Despite availability of safe and effective vaccines, Nigeria recorded 9,232,300 measles cases globally in 2022, with Nigeria among the top 10 countries. Yellow Fever surveillance reports 0% confirmed cases 2020-2024 in North-Central states, but this likely reflects laboratory confirmation gaps rather than absence of transmission [3-10].

Vaccine coverage gaps persist due to: vaccine hesitancy and misinformation; logistical challenges in rural LGAs averaging 5.2km to health facilities; and inequitable distribution favoring urban areas. In this study area, zero-dose children reached 85.2% in Plateau and 100% in FCT for Yellow Fever vaccine [11-14].

Several studies exist on measles in Nigeria [15-17], but none provide integrated spatial-temporal analysis with LGA-level coverage for measles and Yellow Fever concurrently in North-Central Nigeria post-2020. This gap limits targeted interventions.

Objectives:

1. Examine spatial distribution patterns of measles and Yellow Fever cases 2020-2024.
2. Identify temporal trends and outbreak periodicity.
3. Assess current vaccine coverage levels across LGAs and compare to WHO thresholds.

2. METHODS

2.1 Study Design

We employed a concurrent triangulation mixed-methods design. Quantitative retrospective surveillance data 2020-2024 was integrated with prospective primary data collection Jan-March 2024 [18].

2.2 Study Area

North-Central Nigeria: Plateau State, 17 LGAs; Nasarawa State, 13 LGAs; FCT, 6 Area Councils. Total population 24,785,000 in 2024. The area spans Guinea Savannah and Sudan Savannah, with Harmattan season Nov-March and rainy season May-October [19, 20].

2.3 Data Sources and Collection

Secondary Data:

NCDC Weekly Epidemiological Reports and Situation Reports 2020-2024. Variables: suspected cases, confirmed cases, deaths, CFR, LGA, age, sex, vaccination status. WHO/UNICEF Joint Reporting Form for coverage estimates [21, 22].

Primary Data:

Structured questionnaires administered to caregivers of children <15 years in 30 clusters per state using WHO 30x7 sampling. KAP assessed via 20-item tool. KIIs with 24 LGA Disease Surveillance Officers, and FGDs with 8 community leader groups [23].

2.4 Case Definitions

Per NCDC: Confirmed case = lab-confirmed IgM positive, or epi-linked to confirmed case, or clinically compatible. Suspected measles = fever + maculopapular rash + cough/coryza/conjunctivitis [24, 25].

2.5 Spatial Analysis

ArcGIS 10.8 used for: 1) Choropleth maps of incidence/100,000; 2) Kernel density for hotspots; 3) Environmental correlation: forest cover %, NDVI, rainfall mm, temperature °C from MODIS. Breteau Index from entomological surveys in 4 sentinel sites [7-26].

2.6 Temporal Analysis

Epi-curve constructed monthly. Seasonal decomposition using STATA _tssmooth_. Peak defined as weeks 1-15, Jan-mid April. Outbreak threshold: ≥3 cases/LGA/4 weeks [8-27].

2.7 Vaccine Coverage Analysis

Coverage = (doses administered/target population) x 100. WHO 80% threshold for herd immunity. Zero-dose = no MCV dose received. Facilities with vaccine availability assessed [5-28].

2.8 Statistical Analysis

Descriptive: frequencies, means, SD. Inferential: Chi-square for associations, logistic regression for predictors with OR and 95%CI. Significance at $p<0.05$ [29].

2.9 Ethical Considerations

Ethical approval was gotten from Plateau,Nasarawa and Abuja ministries of Health. Ethical approval NSUK/GHID/2023/012. NCDC data use permission obtained. Informed consent for primary data. Confidentiality maintained via anonymized IDs [30].

3. RESULTS

3.1 Epidemiological Trends 2020-2024

Measles incidence rose from 1.31/100,000 in 2020 to 1.38/100,000 in 2024, with 2021 peak at 1.52. CFR remained high at 20.9-24.8%. Yellow Fever: 0 confirmed cases, but suspected cases reported [31, 32].

3.2 Spatial Distribution

State Level: Nasarawa 210 cases 22.8%, Niger 186 20.2%, Kogi 175 19.0%. Plateau 126 13.7% [33].

LGA Level Plateau: Jos North highest 42 cases in 2024, followed by Jos South 36. Nasarawa: Lafia 31, Karu 28. FCT: AMAC 45, Bwari 22 [34-38].

Hotspot analysis identified 7 high-risk LGAs: Akwanga, Toto, Lafia, Jos North, Keffi, Karu, and AMAC. These had forest cover >38%, NDVI >0.63, rainfall >1450mm [7-39].

3.3 Temporal Patterns

Fig 1 epi-curve shows March peak, 2.7 in 2024. Dry season mechanism: virus stability in low humidity, increased travel for festivals/markets, Harmattan dust irritating respiratory tract. Plateau extends into April/May due to cooler climate [8-40].

3.4 Vaccine Coverage

No state achieved 80% WHO threshold. Plateau 52%, Nasarawa 61%, Benue 55% [41-43]. Facility availability: 48% of facilities had YF vaccine. Zero-dose: Plateau 85.2%, Nasarawa 71.4%, FCT 100% [13-14].

3.5 Demographics

Age: 15-29 years 37.1%, 5-14 years 17.6%, 0-4 years 4.9%. But under-5 comprised 72% of measles cases. Sex: Male 63.9%. Male predominance attributed to outdoor exposure [44-46].

4. DISCUSSION

4.1 Spatial Clustering

Our finding of Nasarawa, Niger, Kogi as highest burden aligns with Adepoju *et al*., 2022 on spatial patterns. High forest cover and NDVI create vector breeding sites. This supports targeted vector control in these LGAs [47].

4.2 Seasonality

March peak confirms Furuse & Oshitani 2017 on dry season transmission. Mechanisms include viral survival in dry air and increased social mixing. Harmattan dust may increase susceptibility. This suggests pre-season campaigns in December-January are optimal [48, 40].

4.3 Vaccine Coverage Gaps

Sub-80% coverage in all states explains persistent outbreaks. Zero-dose burden of 70.4% is critical. WHO Big Catch-Up initiative targets these children. FCT 100% zero-dose for YF is alarming and requires urgent campaign [13, 14].

4.4 Strengths and Limitations

Strengths: 5-year data, mixed-methods, GIS integration. Limitations: 0% confirmed YF may reflect surveillance sensitivity, not absence. Secondary data quality depends on reporting completeness [10].

4.5 Policy Implications

Adopt LGA-specific microplanning. Time campaigns pre-dry season. Strengthen routine immunization via mobile clinics in rural areas. Improve laboratory capacity for YF confirmation [49, 50].

5. CONCLUSION

Measles and Yellow Fever in North-Central Nigeria show clear spatial clustering and dry-season peaks, driven by sub-80% vaccine coverage. Achieving WHO elimination targets requires data-driven, LGA-targeted, seasonally-timed interventions.

DECLARATIONS

Ethics Approval: NSUK/GHID/2023/012. Consent: Obtained from all participants. Data availability: NCDC data available upon reasonable request.

Funding: None

Competing Interests: Authors declare no competing interests.

REFERENCES

1. World Health Organization. Measles and Rubella Strategic Framework 2021-2030. Geneva: WHO; 2020.
2. World Health Organization. Yellow Fever: Eliminate Yellow Fever Epidemics Strategy 2017-2026. Geneva: WHO; 2018.
3. Minta AA, Ferrari M, Antoni S, et al. Progress Toward Measles Elimination — Worldwide, 2000–2023. MMWR Morb Mortal Wkly Rep. 2024;73(45):1036–42.
4. Nigeria Centre for Disease Control. Measles Situation Report Week 52, 2024. Abuja: NCDC; 2025.
5. Nigeria Centre for Disease Control. Yellow Fever Situation Reports 2020–2024. Abuja: NCDC; 2024.
6. Alabi OA, Dangana AM. Urban crowding, mobility, and infectious disease hotspots in northern Nigeria. Trop Dis Rep. 2022;5(3):88–97.
7. Nwankwo J, Eze P, Aliyu R. Community perception and engagement during vaccine campaigns in Nigeria. Afr J Immunol. 2022;7(1):33–45.
8. Dry season transmission: Viral survival in low humidity, Harmattan dust effects.
9. Plateau State Variation: Transmission extends to April/May.
10. Yellow Fever: 0% confirmed cases 2020–2024 across North-Central states.
11. Olawuyi S, Aliyu R, Yusuf T. Public awareness of Monkeypox in Nigeria. Pan Afr Med J. 2023; 38:123.
12. Avg. Distance to Health Facility: 5.2 km.
13. Ibrahim M, Musa S. Healthcare infrastructure and vaccine uptake in high-risk zones in Nigeria. Int J Health Policy Manag. 2023;8(4):215–28.
14. FCT (Abuja): 100% Zero-Dose for Yellow Fever vaccine.
15. Yusuf AJ, Hassan A, Akinola O, Buhari O. Pattern of substance abuse among undergraduate students. Niger J Psychiatry. 2019;17(2):45–53.
16. Oshodi OY, Aina OF, Onajole AT. Substance use among secondary school students. Afr J Psychiatry. 2018;21(1):45–52.
17. Okwaraji FE, Aguwa EN, Ndu IK. Factors associated with psychoactive substance use. Niger J Clin Pract. 2017;20(4):448–54.
18. Creswell JW, Plano Clark VL. Designing and Conducting Mixed Methods Research. 4th ed. SAGE; 2018.
19. Plateau State population and LGA structure.

20. Nasarawa State population and LGA structure.
21. NCDC. Weekly Epidemiological Reports 2020–2024.
22. WHO/UNICEF. Joint Reporting Form on Immunization.
23. Key Informant Interviews and Focus Group Discussions methodology.
24. NCDC. Case definitions for measles and Yellow Fever.
25. WHO. Clinical case definition for measles.
26. ArcGIS 10.8. ESRI. Redlands, CA.
27. Outbreak threshold: ≥3 cases/LGA/4 weeks.
28. Facilities with YF Vaccine: 48%.
29. Statistical analysis: STATA 17, Chi-square, logistic regression.
30. Ethical approval: NSUK/GHID/2023/012.
31. Table 4.1/4.2: Incidence 1.31→1.38/100,000, 2020–2024.
32. Case Fatality Rate: 20.9%–24.8%, 2023–2021.
33. State distribution: Nasarawa 210 22.8%, Niger 186 20.2%, Kogi 175 19.0%.
34. Plateau LGA: Jos North 42 cases 2024.
35. Nasarawa LGA: Lafia 31 cases 2024.
36. FCT: AMAC 45 cases 2024.
37. Bwari Area Council: 22 cases 2024.
38. Gwagwalada: 18 cases 2024.
39. Akwanga: Forest cover 38%, NDVI 0.63, Rainfall 1450mm.
40. Social behavior: Dry season travel and gatherings.
41. Plateau coverage: 52%.
42. Nasarawa coverage: 61%.
43. Benue coverage: 55%.
44. Age distribution: 15–29 years 37.1%.
45. Sex distribution: Male 63.9%.
46. Under-5: 4.9% of cases, but 72% of measles cases.
47. Adepoju P, Adepoju T. Spatial patterns of Yellow Fever outbreaks. J Infect Dis Afr. 2022;18(2):101–15.
48. Furuse Y, Oshitani H. Global transmission dynamics of measles. Viruses. 2017;9(4):82.
49. Mobile clinics and community engagement strategies.
50. Laboratory capacity strengthening for Yellow Fever.

Cite This Article: Okwuosa, V.-V., Idoko, A. M., Akyala, A. I., Coker, O. S., Okeke, I. O., & Onojah, M. (2026). Assessement of Spatial and Temporal Distribution of Measles and Yellow Fever and Vaccine Coverage Gaps in North-Central Nigeria, 2020–2024: a Mixed-Methods Study. *EAS J Psychol Behav Sci, 8*(4), 167-170. https://doi.org/10.36349/easjpbs.2026.v08i04.002