

Original Research Article

Biochemical Characterization and Amino Acid Profile of some Local Resources Used in Food Supplementation

Aminata Sanou¹, François Tapsoba¹, Jeanne d'Arc Wendmintiri Kabré¹, Sanogo Bougma¹, Mah Alima Esther Traoré², Prudence Dounia¹, Charles Parkouda², Aly Savadogo^{1*}

¹Laboratoire de Biochimie et Immunologie Appliquées (LaBIA), Département de Biochimie-Microbiologie, Université Joseph KI-ZERBO, 03 BP7021 Ouagadougou 03, Burkina Faso

²Département de Technologie Alimentaire, Institut de Recherche en Sciences Appliquées et Technologies (DTA/IRSAT), 03 BP 7047, Ouagadougou 03, Burkina Faso

Article History

Received: 15.07.2023

Accepted: 26.08.2023

Published: 31.08.2023

Journal homepage:

<https://www.easpublisher.com>

Quick Response Code



Abstract: Nutritional studies make little reference to essential amino acids to explain undernutrition in protein-energy malnutrition. This study aimed to carry out biochemical characterization and amino acid profiling of *Detarium microcarpum* and *Adansonia digitata* fruit pulps and *Moringa oleifera* leaf powder to show their contribution to protein balance. Biochemical analyses and amino acid profiles were carried out using standard methods and statistically analyzed using XLstat 2016 software. Results are expressed on a dry matter basis. Moringa powder had the highest ash, lipid and protein contents, at 10.72%, 8.75% and 23.85% respectively, compared with 4.14%, 1.99% and 4.19% for *Detarium*. *Adansonia* pulp had the lowest lipid and protein contents, at 1.36% and 1.83% respectively. In terms of total amino acids, Moringa powder had the highest content at 15.59%, while baobab pulp had the lowest at 2.18%. Essential amino acids for Moringa powder were higher and more balanced than for fruit. Histidine was absent in *Adansonia digitata* pulp, while methionine was absent in both *Adansonia digitata* and *Detarium microcarpum* pulps. Thus, blending the three resources could result in products with high nutritional value.

Keyword: *Adansonia digitata*, Amino acids, *Detarium microcarpum*, Food supplementation, *Moringa oleifera*, Nutritional composition.

Copyright © 2023 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution **4.0 International License (CC BY-NC 4.0)** which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

Plants have always been an important source of nutrition for many living beings. In some parts of the world, due to health problems, dietary choices or economic reasons, plant products are the only way to obtain all the nutrients the body needs especially when resources are very limited (Burlandoa *et al.*, 2019). In the Sahelian landscape, the three plants *Detarium microcarpum* (sweet detar), *Adansonia digitata* (Baobab) and *Moringa oleifera* are of particular importance in terms of diet and therapeutic practices. In rural areas, women and children are responsible for harvesting the fruits and leaves of these plants (MEDD, 2012; Ricci *et al.*, 2020). Their importance in nutrition and traditional medicine is based on their mineral and vitamin content, their antioxidant power and the presence of numerous other biological molecules of therapeutic aptitude. Indeed, in 2019, a study presented the pharmacological and therapeutic properties of

Detarium microcarpum, including antioxidant, anti-diabetic, antibacterial, anti-fungal, antiviral and anti-cancer activities, as well as its nutritional and functional properties (Burlandoa *et al.*, 2019). Monkey bread, the fruit of *Adansonia digitata*, has been shown to help maintain phospho-calcium balance, in addition to its acidic nature due to its organic acids (Parkouda *et al.*, 2007). *Moringa oleifera* powder has long been known for its qualities in the nutritional recovery of immunocompromised people, notably following infection by the Human Immunodeficiency Virus (HIV), in preventing lipid oxidation and other food-altering reactions (Hodas *et al.*, 2021) and in reducing aflatoxin B1 levels (Gard *et al.*, 2021). Its virtues are such that Moringa is called 'arzantiga' or tree of paradise in the local Moore language. Unfortunately, the dietary benefits of Moringa are negatively influenced by its low consumption (Zongo *et al.*, 2013). In Benin, a 20% refusal rate by malnourished children

*Corresponding Author: Aly Savadogo

Laboratoire de Biochimie et Immunologie Appliquées (LaBIA), Département de Biochimie-Microbiologie, Université Joseph KI-ZERBO, 03 BP7021 Ouagadougou 03, Burkina Faso

aged 6 to 59 months was recorded by researchers following the administration of a Moringa-enriched diet due to taste and smell (Agossadou *et al.*, 2018).

On the other hand, protein requirements boil down to the amino acids needed to synthesize body proteins and other nitrogenous substances (Kenfack, 2010). In nutrition, the notion of protein quality is thus relative to its digestibility and amino acid composition. While researchers agree that digestibility depends on the presence of several factors in the feed, some of which are unfavorable (anti-nutritional factors such as tryptic inhibitors), amino acids in turn are assessed based on their essential amino acid (EAA) content. In the absence of essential amino acids, the body is unable to synthesize specific proteins (Darmaun, 2008). Yet (Müller and Krawinkel, 2005), proteins are irreplaceable by other nutrients. This results in a low protein mass following a negative nitrogen balance, with consequences such as undernutrition and its attendant morbidities (Hsairi *et al.*, 2021). Studies have extolled the virtues of Moringa, Detarium and monkey bread as supplements, particularly for vulnerable people, basing their approach on micronutrients (minerals, trace elements and vitamins). However, the investigation of amino acids as essential micronutrients for maintaining healthy body function has often remained a secondary priority. The present study is therefore a contribution to our knowledge of the essential amino acid composition of products from three Sahelian plants, to clarify their interest in supplementation. The study aimed to characterize the biochemistry and amino acid profile of *Detarium microcarpum* and *Adansonia digitata* fruit pulps and *Moringa oleifera* leaf powder used for food fortification in Africa.

MATERIAL AND METHODS

Obtaining plant material

The plant material consisted of *Detarium microcarpum* fruits, *Adansonia digitata* fruits and *Moringa oleifera* leaves. *Detarium microcarpum* and *Adansonia digitata* fruits were obtained from women growers in the villages surrounding the town of Gaoua. The fruits were packed in cartons and transported at room temperature to the food technology workshop. Moringa leaves were obtained from the vegetable garden in the town of Gaoua. The freshly picked leaves were packed in bags and transported to the food technology workshop for drying and grinding into powder.

Production of fruit pulp and dried leaf powder

Detarium microcarpum fruits, *Adansonia digitata* hulls (monkey bread) and *Moringa oleifera*

leaves were processed according to the diagrams shown in Figure 1.

Pre-treatment of *Detarium* fruit involves sorting, washing and drying. These steps enabled us to eliminate fruit with defects, skins adhering intimately to the pulp, sand and surface dirt, and to make the skins drier. Processing consists of extracting the powder by crushing, followed by sieving to retain almond fibers and debris. The resulting pulp is then packaged for consumption.

A series of operations, including brushing and breaking the shell, removed the yellowish-green layer covering the shell and extracted the contents, which consisted of the *Adansonia digitata* pulp surrounding a kernel, all entangled in a network of fibers. The dry pulp was obtained by crushing and sieving.

The moringa powder production protocol was developed by (Sauveur and Broin, 2010). The treatment applied to moringa petioles consisted in stripping the leaves, washing them in a 1% saline solution for around 3 minutes before draining them and then drying them at 50°C for at least 6 hours in an electric dryer. The dried leaves were ground and sieved to produce Moringa powder, which was packaged in food-safe containers.

Determination of physicochemical and nutritional characteristics of dried leaf pulp and powder

Physicochemical and nutritional characteristics of dried leaf pulp and powder were determined using standard methods.

Water content was determined by weight difference before and after oven drying at 105°C to a constant weight (AOAC, 2000).

Ash was determined using the ISO 2171 method by incinerating samples in a muffle furnace at 550°C. The ash was cooled in a desiccator and weighed. (ISO, 2007).

The Kjeldahl method (AFNOR, 1986) was used for nitrogen determination. The nitrogen determined after mineralization followed by distillation is multiplied by the conversion factor 6.25.

Lipids were extracted in hexane using the Soxhlet followed by oven drying according to (AOAC, 2000).

Fibers were determined after digestion with sulfuric acid and neutralization with sodium hydroxide under boiling conditions, washing with hot water, filtration, oven-drying at 105°C for 8 hours, incineration in a furnace at 550°C for 3 hours according to the method (AOAC, 1990).




Figure 1: *Detarium microcarpum* pulp (a), *Adansonia digitata* pulp (b) and *Moringa oleifera* leaf powder (c)

Digestible carbohydrates were determined by the differential method according to the formula proposed by (Egan *et al.*, 1981). Digestible carbohydrate content (%) = 100 - [water content (%) + protein content (%) + lipid content (%) + ash content (%) + fiber content (%)].

Energy value was obtained by multiplying macromolecule content by Atwater's coefficients: VA (kcal/100g) = [(% Carbohydrates x 4) + (% Protein x 4) + (% Fat x 9)] (Atwater and Benedict, 1899).

Amino acids were determined by HPLC using the PICO-TAG method developed by the DANIDA Project of the Danish Technological Institute (Kristoffersen, 2011).

Statistical analysis of data

Data were entered into Microsoft Excel 2016 and compared using ANOVA to assess differences between means. The Fisher test was used to compare variances at the 5% significance level. A Principal Component Analysis (PCA) using XLSTAT version

2016 was performed to show the correlation between essential amino acids and plant resources.

RESULTS

Biochemical characteristics and energy value of pulp and leaf powder

The physicochemical characteristics of *Detarium microcarpum* pulp, *Adansonia digitata* and *Moringa oleifera* leaf powder are summarized in Table 1.

Adansonia digitata pulp had a higher moisture content than *Detarium microcarpum* pulp and *Moringa oleifera* powder. *Moringa oleifera* is a good source of protein and lipids, while pulp is an excellent carbohydrate reserve. *Detarium microcarpum* fruit pulp showed a higher energy value, while Moringa powder revealed an ash content twice that of *Adansonia digitata* fruit pulp. For all physicochemical parameters, the difference was statistically significant at $p < 0.01$. See below.

Table 1: Physicochemical and nutritional parameters and energy value of *Detarium* and baobab fruit pulps and *Moringa* powder expressed as dry matter.

Composition	Detarium Pulp	Adandonia Pulp	Moringa powder	P-value
Humidity (%)	6.59±0.17	9.15±0.09	6.75±0.17	< 0.01
Ashes (%)	4.14±0.12	5.87±0.33	10.72±0.00	< 0.01
Lipids (%)	1.99±0.05	1.36±0.08	8.75±0.4	< 0.01
Proteins (%)	4.19±0.13	1.83±0.13	23.85±0.63	< 0.01
Fibers (%)	8.44±0.07	9.98±0.11	15.40±1.40	< 0.01
Digestible carbohydrates (%)	74.64±0.00	71.81±0.16	34.51±0.37	< 0.01
Energy value (kcal/100g)	333.30±0.05	306.81±0.85	312.86±7.62	< 0.01

For moisture, protein, lipids, fiber and digestible carbohydrates, values are averages of two determinations $\pm$ standard deviation. For ash, values are averages of three determinations $\pm$ standard deviation.

Total amino acid content of Detarium pulp, *Adansonia digitata* and Moringa leaf powder

A total of 17 amino acids were used as standards. The proportion of total amino acids in Detarium and *Adansonia digitata* pulps and Moringa leaf powder is shown in the histogram in Figure 2. Moringa powder contained more amino acids than Detarium pulp, which contained more than *Adansonia digitata* pulp.




Figure 2: Total amino acid content of *Detarium microcarpum* pulp, *Adansonia digitata* pulp and *Moringa oleifera* powder

Principal component analysis of essential amino acids

Principal component analysis (Fig 3) was used to establish the correlation biplot between the essential amino acids quantified in the three local products: *Detarium microcarpum* and *Adansonia digitata* pulps and *Moringa oleifera* leaf powder. The first two factorial axes F1/F2 show 100% variability in eigenvalues; this means that the representation is of good quality. Threonine, alanine, valine, methionine,

isoleucine, leucine, phenylalanine and histidine were collinear and significantly positively correlated with lysine. In addition, the F1 axis was more closely related to these variables, while both the F1 and F2 axes were equidistant from lysine. In terms of observations, Moringa powder was almost conflated with the F1 axis, while *Detarium microcarpum* and *Adansonia digitata* pulps had a more nuanced contribution. These two resources shared the subspaces formed by the intersection of the F1 and F2 dimensions see below.




Figure 3: F1/F2 axis correlation biplot of essential amino acids

DISCUSSION

The moisture content of *Adansonia digitata* pulp was higher than that indicated in NBF 01-213, i.e. 8.7% (ABNORM, 2018a). This could be because the pulp extracted from the epicarp did not undergo drying. However, this content was close to that found by (Tapsoba *et al.*, 2023), which was 9.63%. In contrast, *Moringa oleifera* powder had a moisture content in line with NBF 01-216, i.e. below 8% (ABNORM, 2018b). Detarium pulp moisture was higher than that reported by a team of researchers from the University of Ouagadougou, i.e. 4.93% (Thiombiano *et al.*, 2014) but lower than the 7.15%, 7.12% and 7.26% respectively at the Bongor, Gaya and Lāi sites (Makalao *et al.*, 2015). The moisture content of Moringa leaf powder was lower than the 8.25% obtained in another study (Ouédraogo *et al.*, 2018). The moisture contents obtained for the fruit pulps indicate that they should be dried further to ensure a long shelf life without spoilage for year-round consumption needs.

Moringa ash content was higher than that obtained from *Adansonia* fruit pulp and corroborates the results reported by other authors. The ash content of *Adansonia digitata* fruit pulp was higher than the average reported by a group of researchers, i.e. 4.9%. (Stadlmayr *et al.*, 2013). Edaphic factors could explain this difference. In addition, for *Detarium microcarpum* pulp, ash contents obtained in the present study were similar to those reported by authors in Cameroon with rates of 4.14% (Bayang *et al.*, 2021). Ash is a nutritional quality criterion as it is the inorganic residue of the mineral-containing feed (Nielsen, 2017). Thus, the present work confirms the nutritious nature of local products often used in improving the nutritional status of vulnerable people.

The fat content of Moringa powder was high compared with that of *Detarium microcarpum* and *Adansonia digitata* fruit pulps. A fat content of 7.85%, lower than in this study, was obtained in Senegal (Ndong *et al.*, 2007). On the other hand, a group of researchers (Pamba *et al.*, 2018) found for *Adansonia digitata* fruit pulp sold on the market in Côte d'Ivoire, fats contents of 0.42 g/100g dry matter, lower than those obtained in this study. However, the results of the present study are close to those of other authors, who found 2.23% on a dry basis (Oibiopka *et al.*, 2014). More generally, Muthai and colleagues had shown that proximal compositions of plant resources differ according to geographical provenance (Muthai *et al.*, 2017). Fats are necessary for the transport of fat-soluble vitamins and enter into the constitution of biological limbs.

The protein content of *Adansonia digitata* fruit pulp was lower than that indicated in the Burkinabe standard, i.e. 2.7% (ABNORM, 2018a), while that of *Detarium* fruit pulp was close to that (Thiombiano *et al.*, 2014), i.e. 4.65%. The protein content of Moringa

leaf powder was also close to that obtained in Mali by Coulibaly's team, i.e. 24.2% (Coulibaly *et al.*, 2022). The age of the plants could be a determining factor in quantitative composition. According to some authors, proteins are more abundant in the leaves of young plants than in those of adult plants (Kane *et al.*, 2010). In addition, dieticians (Pamplona-Roger, 2006) assert that it is easier to ingest large quantities of fruit than other foods such as meat. So, although overall protein levels in the fruits studied were lower, in the lean season when they are generally consumed a lot, their contribution to dietary balance is not negligible, given the inaccessibility of animal proteins to poor households (Kenfack, 2010).

The fiber content of *Detarium* fruit pulp was higher than the 7.87% on a dry basis (Jacob *et al.*, 2016), while the content for baobab pulp was lower than the results of (Compaoré *et al.*, 2011) i.e. 11.87%. These disparities could be linked to the maturity stage of the parties involved. Furthermore, (Cissé, 2012) finds an explanation for these differences in the diversity of methods used for the same biological material. The fiber value of Moringa powder was close to that found by (Yaméogo *et al.*, 2011) i.e. 15.7%.

The digestible carbohydrate values of baobab fruit pulp were close to those of (Pamba *et al.*, 2018), i.e. 71.39% for the Yopougon market. In contrast, the content of *Detarium* pulp obtained in this study was higher than that obtained in Nigeria, i.e. 65.38% digestible carbohydrates (Oibiopka *et al.*, 2014). These disparities could be explained by climatic conditions or the degree of ripeness of the fruit, among other factors. In fact, according to some authors, fruit sugar content and acidity evolve inversely: at maturity, acids tend to decrease while sugar content increases (Tyl and Sadler, 2017). Given the high digestible carbohydrate content of *Detarium microcarpum* fruit pulp, it can be used as a sugar substitute in Sudan (Cavin, 2007). This ability of *Detarium microcarpum* fruit pulp could be used in conjunction with its other attributes to enrich infant flours.

The energy value of baobab pulp was much higher than those reported by (Makalao *et al.*, 2015) i.e. from 65.46 kcal/100g to 68.1 kcal/100g but similar to those (Pamba *et al.*, 2018) i.e. 306.08 kcal/100g for the Yopougon market. Compared with samples from other authors (Thiombiano *et al.*, 2014), the values obtained in this study are close for *Detarium* fruit pulp (i.e. 335.5 kcal/100g). Moringa powder presented a lower energy value than that found in Mali, i.e. 396 kcal for 100g of dry Moringa leaf powder (Coulibaly *et al.*, 2022). This difference would be due to the quantitative difference in calorific molecules (carbohydrates, lipids, proteins). Thus, each of these resources can contribute to improving energy density as part of a supplemental diet. In addition, the combination of *Detarium microcarpum* and *Adansonia digitata* fruit pulps and Moringa powder

constitutes a concentrate of protein-energy nutrients. Such a compound would be effective in the fight against recurrent protein-infant malnutrition in Sub-Saharan Africa, resulting from a diet whose energy density is often deemed low about energy needs of young children.

In terms of overall amino acid composition, Moringa powder showed the highest content compared to Detarium and Adansonia fruit pulps. Total amino acids, 26.032%, had been identified by Moyo and colleagues in dried Moringa leaves (Moyo *et al.*, 2011). These levels were higher than in this study. This could be explained by edapho-climatic factors, the difference in variety or the number of amino acids. Indeed, Moyo's team had determined nineteen (19 amino acids), whereas we had only looked for seventeen (17). Furthermore, in terms of quality, Moringa powder contains the eight essential amino acids in a more balanced proportion. These results shed further light on Moringa powder as a resource for supplementing protein undernutrition in Africa. When comparing Malagasy and Ivorian baobab pulps, (Cissé, 2012) found six (6) free amino acids in the Ivory Coast species *Adansonia digitata*, whereas we found fifteen (15). Among others, we found lysine and leucine, which were only detected in certain Madagascar species by the same author. Furthermore, (Cissé, 2012) had reported the lowest values for the sulphur amino acids cysteine and methionine, whereas the present study revealed an absence of methionine but a cysteine content that exceeds that of valine, tyrosine, alanine, glycine, serine and glutamic acid. Methionine was not detected in Detarium pulp either. Thus, the protein contribution of pulps in essential amino acids is only partial. In his book on food science, Pamplona-Georges (Pamplona-Roger, 2006) revealed the absence of methionine among fruit amino acids, which corroborates the results of this study. Consequently, this author asserted that the amino acids in fruit are incomplete compared to those in vegetables. So, for a balanced diet, you need both fruit and vegetables. Hence the need to combine Detarium and Adansonia fruit pulps with Moringa powder for effective supplementation of essential amino acids. We plan to supplement infant flours with a combination of these three products to assess their essential amino acid profile.

CONCLUSION

Moringa oleifera leaves and *Detarium microcarpum* and *Adansonia digitata* pulps are nutrient reserves in terms of their macromolecular and micronutrient composition. Moringa powder has a higher nutritional value than pulp, particularly in terms of total and essential amino acids. The combination of Detarium and Adansonia pulps increases nutritional composition. However, the absence of methionine in each of the pulps makes the quality of their proteins low. For supplementation needs, we recommend

combining *Moringa oleifera* powder with *Adansonia digitata* pulp and/or *Detarium microcarpum* pulp.

ACKNOWLEDGMENTS

The authors are grateful to the Department of Food Technology of the Research Institute of Applied Sciences and Technologies (DTA/IRSAT) and the Laboratory of Applied Biochemistry and Immunology (LaBIA) for providing the platform and reagents for the analysis.

Funding Sources: The authors received no financial support for the research, authorship, and/or publication of this article.

Conflicts of interest: The authors declare no conflict of interest.

Consent to publication

All the authors confirm the manuscript have not been published elsewhere, have not been considered by any other publisher. The all consent to the publication of the manuscript in EAS Journal of Nutrition and Food Sciences.

REFERENCES

- ABNORM. (2018a). Norme burkinabè nbf 01-213: 2018_ pulpe de fruit de baobab (pain de singe): Spécifications.
- ABNORM. (2018b). Norme burkinabè nbf 01-216: 2018_ poudre de feuilles de moringa: Spécifications.
- AFNOR. (1986). Directives générales pour le dosage de l'azote avec minéralisation selon la méthode de kjedahl. Produits agricoles alimentaires NF V 03-050-1970.
- Agossadou, J. O., Houndji, B. V. S., Somessi, A. B., Nago, E. S. H., & Chadare, F. J. (2018). Effectiveness of cochlospermum tinctorium root powder based sauce fortified with moringa leaf powder in improving nutritional status of children aged 6 to 59 months in northern benin, 17(2), 352-357.
- AOAC. (1990). Official methods of analysis_ agricultural chemicals; contaminants, drugs. Edited by Kenneth Helrich Fifteenth Edition, 1, 80-82.
- AOAC. (2000). Official methods of analysis. Association of official analytical chemists, Washington D.C., USA.
- Atwater, W. O., & Benedict, F. G. (1899). Experiments on the metabolism of matter energy in the human body. US Department of Agriculture Washington, DC Bulletin, 69, 112.
- Bayang, J. P., Laya, A., Kolla, M. C., & Koubala, B. B. (2021). Variation of physical properties, nutritional value and bioactive nutrients in dry and fresh wild edible fruits of twenty-three species from Far North region of Cameroon. *Journal of Agriculture and Food Research*, 4, 100146.

- Burlando, B., Palmero, S., & Cornara, L. (2019). Nutritional and medicinal properties of underexploited legume trees from West Africa. *Critical reviews in food science and nutrition*, 59(sup1), S178-S188.
- Cavin, A. L. (2007). Contribution à la connaissance taxonomique et chimique de fruits africains du genre "detarium" (fabaceae - caesalpinioideae): "D. Microcarpum" guill. Et perr. Et des formes comestibles et toxiques de "d. Senegalense" j.F. Gmel. Thèse de doctorat: Univ. Genève, 2007, no. Sc. 3838(p.42).
- Cissé, I. (2012). Caractérisation des propriétés biochimiques et nutritionnelles de la pulpe de baobab des espèces endémiques de madagascar et d'afrigue continentale en vue de leur valorisation. Université de Montpellier Supagro.
- Compaoré, W. R., Nikiéma, P. A., Bassolé, H. I. N., Savadogo, A., & Mouecoucou, J. (2011). Chemical composition and antioxidative properties of seeds of Moringa oleifera and pulps of Parkia biglobosa and Adansonia digitata commonly used in food fortification in Burkina Faso. *Current Research Journal of Biological Sciences*, 3(1), 64-72.
- Coulibaly D., Dembélé Y. K., Diawara F., Touré O. B., Iknane A.A., & Sanogo R. (2022). Caractérisation et composition biochimique et nutritionnelle du Moringa Oleifera récolté dans le district de Bamako, Mali en 2020. *Mali Santé Publique*, 5-13.
- Darmaun, D. (2008). Qu'est-ce qu'un acide aminé essentiel en 2008? *Nutrition clinique et métabolisme*, 22, 142-150.
- Egan, H., Kirk, R. S., & Sawyer, R. (1981). Pearson's chemical analysis of food. (8th edition). Churchill. Livingstone: London-UK:pp 591.
- Gard, C., Kolich Chong, M. A., Sawadogo, Y. A., & Moreau, L. (2021). Plant addition impact on aflatoxin B1 levels, physico-chemical, nutritional, and sensory properties of fried peanut biscuits. *Journal of Food Safety*, 41(3), e12894.
- Hodas, F., Zorzenon, M. R. T., & Milani, P. G. (2021). Moringa oleifera potential as a functional food and a natural food additive: A biochemical approach. *Anais da Academia Brasileira de Ciencias*, 93(Suppl. 4), e20210571.
- Hsairi, M., Gargouri, L., Safi, F., Elleuch, A., & Mahfoudh, E. (2021). Denutrition Proteino-Energetique Chez L'enfant Protein Energy Malnutrition In Children. *Journal de l'Information Médicale de Sfax*, 37, 10-16.
- ISO. (2007). Céréales, légumineuses et produits dérivés_ dosage du taux de cendres par incinération. Fourth edition 2007-11-01, 1-9.
- Kane, F., Yao-Kouamé, A., Konan, A. A., & N'Guessan, A. K. (2010). Dosage de quelques composantes biochimiques des feuilles de lippia multiflora (verbenacee) à deux stades de développement et qualité des infusions en fonction de la dose d'urée. *Agronomie Africaine*, 22(3), 227-235.
- Kenfack, L. B. M. (2010). Propriétés nutritionnelles et fonctionnelles des protéines de tourteaux, de concentrats et d'isolats de *ricinodendron heudelotii* (bail.) pierre ex pax et de *tetracarpidium conophorum* (müll. Arg) [Thèse]. Université de Ngaoundéré.
- Kristoffersen, L. T. (2011). Amino acid analysis of savannah tree seeds. Danish Technological Institute.
- Makalao, M. M., Savadogo, A., Zongo, C., & Traore, A. S. (2015). Composition nutritionnelle de 10 fruits sauvages consommés dans trois départements du tchad International Journal of Biological and Chemical Sciences-IJBCS. *Int J Biol Chem Sci*, 9(5), 2385-2400.
- MEDD. (2012). Stratégie nationale de promotion et de valorisation des produits forestiers non ligneux (snpv/pfnl). pp 8.
- Moyo, B., Masika, P. J., Hugo, A., & Muchenje, V. (2011). Nutritional characterization of moringa (moringa oleifera lam.) leaves. *African Journal of Biotechnology*, 10(60), 12925-12933.
- Müller, O., & Krawinkel, M. (2005). Malnutrition and health in developing countries. *CMAJ* 2, 173(3), 279-286.
- Muthai, K. U., Karori, M. S., Muchugi, A., Indieka, A. S., Dembele, C., Mng'omba, S., & Jamnadass, R. (2017). Nutritional variation in baobab (Adansonia digitata L.) fruit pulp and seeds based on Africa geographical regions. *Food science & nutrition*, 5(6), 1116-1129.
- Ndong, M., Wade, S., Dossou, N., Guiro, A. T., & Gning, R. D. (2007). Valeur nutritionnelle du *moringa oleifera*, étude de la biodisponibilité du fer, effet de l'enrichissement de divers plats traditionnels sénégalais avec la poudre des feuilles. *African Journal of Food Agriculture Nutrition and Development (AJFAND)*, 7(3).
- Nielsen, S. S. (2017). Food analysis laboratory manual. Springer Food Science Text Series. Volume Third Edition.
- Oibiokpa, F. I., Saidu, A. N., & Shittu, K. O. (2014). Nutritional composition of Detarium microcarpum fruit. *African Journal of Food Science, academic Journal*, 8(6), 342-350.
- Ouedraogo, S., Traoré, S., Sombié, B. C., Traoré, T. K., Nitiéma, M., BeLeMNaBa, L., ... & GUISSOU, I. (2018). Contrôle qualité et étude de comprimabilité des poudres de feuille de Moringa oleifera (lam) et de pulpe du fruit de Adansonia digitata: quality control and compressibility study of Moringa oleifera powders and Adansonia digitata fruit pulp. *Sciences de la Santé*, 41(2), 97-99.
- Pamba, K. B. A., Assanvo, J. B., & Koffi, K. M. (2018). Caractérisation biochimique et microbiologique de la pulpe de baobab (*adansonia digitata*) vendue sur le marché d'abidjan. *American Journal of Innovative Research and Applied Sciences*.
- Pamplona-Roger, G. D. (2006). Guide des aliments et leur pouvoir curatif. (Editorial Safeliz, S.L).
- Parkouda, C., Diawara, B., Ganou, L., & Lamien, N. (2007). Potentialités nutritionnelles des produits de 16 espèces fruitières locales aux burkina faso. *Science et technique*, 1, pp 35-47.

- Ricci, E., Kaboré, P., Emmenegger, R., & Kaboré, O. (2020). Capitalisation d'expériences de 12 ans de projets au burkina faso. Retour d'expériences dans la valorisation de produits forestiers non ligneux (pfnl). Processus de développement et acquis. ADAP G, Suisse, editor.
- Sauveur, A. D. S., & Broin, M. (2010). Produire et transformer les feuilles de moringa. pp 35-41.
- Stadlmayr, B., Charrondiere, U. R., Eisenwagen, S., Jamnadass, R., & Kehlenbeck, K. (2013). Nutrient composition of selected indigenous fruits from sub-saharan africa. *Journal of Science Food Agriculture (JSFA)*, 93, 2627–2636.
- Tapsoba, F., Zio, S., Tarnagda, B., Waongo, B., Zongo, C., Savadogo, A. (2023). Biochemical, nutritional and microbiological quality of baobab (*adansonia digitata* l.) fruit pulp marketed in ouagadougou. *International Journal of Biochemistry Research & Review*, 32, 1-10.
- Thiombiano D. N. E., Parkouda C., Lamien N., Séré A., Castro-Euler A. M., & Boussim I. J. (2014). Nutritional composition of five food trees species products used in human diet during food shortage period in Burkina Faso. *African Journal of Biotechnology*, 13(17), 1807-1812.
- Tyl, C., & Sadler, G. D. (2017). pH and titratable acidity. *Food analysis*, 389-406.
- Yaméogo, C. W., Bengaly, M. D., Savadogo, A., Nikiema, P. A., & Traore, S. A. (2011). Determination of chemical composition and nutritional values of *Moringa oleifera* leaves. *Pakistan journal of nutrition*, 10(3), 264-268.
- Zongo, U., Savadogo, A., Zoungrana, S. L., Sekoné, P. L., & Traoré, A. (2013). Interêt nutritionnel de *moringa oleifera* lam (syn. *Moringa pterygosperma* c.F. Gaertn.) *Ethnopharmacologia*, 50, p. 36.

Cite This Article: Aminata Sanou, François Tapsoba, Jeanne d’Arc Wendmintiri Kabré, Sanogo Bougma, Mah Alima Esther Traoré, Prudence Dounia, Charles Parkouda, Aly Savadogo (2023). Biochemical Characterization and Amino Acid Profile of some Local Resources Used in Food Supplementation. *EAS J Nutr Food Sci*, 5(4), 119-126.
